

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

1. REPORT DATE (DD-MM-YYYY) 02-16-2011		2. REPORT TYPE Conference Proceeding		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Observations of Higher-Frequency Variability Through the Turkish Straits System				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 0601153N	
				5d. PROJECT NUMBER	
6. AUTHOR(S) Jeffrey W. Book, Ewa Jarosz and Sukru Besiktepe				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER 73-9441-09-5	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7330--09-0098	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660				10. SPONSOR/MONITOR'S ACRONYM(S) ONR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT From September 2008 through February 2009, Acoustic Doppler Current Profilers (ADCPs) measured flow in pairs at the four entrances and exits to the Turkish Strait System. Barotropic tide structures were observed for both diurnal and semidiurnal frequencies in the Aegean/Dardanelles entrance. At the Dardanelles/Sea of Marmara entrance some baroclinic structure was observed with diurnal tides having higher energy in the upper layer compared to the lower layer. Baroclinic structure in the semi-diurnal tides was observed at the Bosphorus/Sea of Marmara entrance and diurnal oscillations were weak. Both diurnal and semi-diurnal oscillations were weak at the Bosphorus/Black sea entrance as the temporal evolution of the flow was more concentrated at low frequencies in the form of a mid-water-column jet.					
15. SUBJECT TERMS Aegean Sea, Black Sea, bosphorus, dardanelles, tides					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18. NUMBER OF PAGES 1	19a. NAME OF RESPONSIBLE PERSON Jeffrey Book
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 228-688-5251

20110224051

OBSERVATIONS OF HIGHER-FREQUENCY VARIABILITY THROUGH THE TURKISH STRAITS SYSTEM

Jeffrey W. Book^{1*}, Ewa Jarosz¹ and Sükrü Besiktepe²

¹ U.S. Naval Research Laboratory, Stennis Space Center, MS, U.S.A. - book@nrlssc.navy.mil

² NATO Undersea Research Centre, La Spezia, Italy

Abstract

From September 2008 through February 2009, Acoustic Doppler Current Profilers (ADCPs) measured flow in pairs at the four entrances and exits to the Turkish Strait System. Barotropic tide structures were observed for both diurnal and semidiurnal frequencies in the Aegean/ Dardanelles entrance. At the Dardanelles/Sea of Marmara entrance some baroclinic structure was observed with diurnal tides having higher energy in the upper layer compared to the lower layer. Baroclinic structure in the semi-diurnal tides was observed at the Bosphorus/Sea of Marmara entrance and diurnal oscillations were weak. Both diurnal and semi-diurnal oscillations were weak at the Bosphorus/Black Sea entrance as the temporal evolution of the flow was more concentrated at low frequencies in the form of a mid-water-column jet.

Keywords: Aegean Sea, Black Sea, Bosphorus, Dardanelles, Tides

The U.S. Naval Research Laboratory in collaboration with NATO Undersea Research Centre and the Turkish Navy Office of Navigation, Hydrography and Oceanography, undertook a scientific research program, Exchange Processes in Ocean Straits (EPOS) that included an observational effort from September 2008 through February 2009 of maintaining pairs of bottom-mounted, upward-looking Acoustic Doppler Current Profilers near all entrances (exits) of the Turkish Strait System. Simultaneous current measurements allow connectivity ranging from high-frequency to seasonal processes to be investigated. One aspect of EPOS was to examine the frequency response of this system which includes very narrow passages and two vertical layers of opposed and energetic flow and exchange [1]. E.g., how do processes that are forced synoptically or faster propagate through the system and what is the baroclinic structure of their evolution?

To investigate this, rotary spectra were calculated from all the ADCP records separately for all depths. We used Welch's averaged periodogram method over block lengths of 43 days and 50% overlapping Hanning windows. No significant polarity differences were found between clockwise and counterclockwise spectra suggesting the absence of inertial activity at all four sections. There is a remarkable difference in the spectra results for each ADCP current section, particularly with regard to tidal oscillations. Diurnal and semidiurnal tides were clearly present at both ends of the Dardanelles Strait. The structures were primarily barotropic at the Aegean/Dardanelles entrance (Fig. 1, left). In contrast, at the Dardanelles/Sea of Marmara entrance the diurnal fluctuations in the lower layer entrance were notably weaker than fluctuations in the upper layer (Fig. 1, right). This suggests that the Dardanelles Strait geometry and/or its baroclinic structure has acted to extract some energy from the barotropic tide and convert it into baroclinic processes. Also, the differences between the diurnal and semidiurnal responses show a frequency dependence on this action.

At the Bosphorus/Sea of Marmara entrance the tidal frequency spectral peaks were different than those observed at the Dardanelles/Sea of Marmara entrance. Diurnal fluctuations were mostly absent and the semi-diurnal fluctuations had baroclinic structure, again with stronger spectral peaks in the upper than the lower layer (Fig. 2, left). This suggests that oscillations at diurnal and semi-diurnal frequency are not passing unaltered through the Sea of Marmara. The observed spectra at the Bosphorus/Black Sea entrance differed the most from the others as spectral energy was concentrated in the middle of the water column and at lower frequencies (Fig. 2, right). Flow was maximized in the upper portions of the lower layer in a jet-like structure as compared to maximums in the upper portions of the upper layer at other sections. Weak (primarily semidiurnal) tidal oscillations persisted in the upper layer and the flow maximum jet, but were virtually absent in the bottom of the water column. This suggests that the Bosphorus Strait geometry and/or its baroclinic structure also acts to alter or block higher frequency oscillations.

Analysis of these results has just begun and is ongoing. However, we can conclude that diurnal and higher frequency oscillations are an important component of the variability of the flow in parts of the Turkish Straits system, and that the Straits System acts to significantly affect the transmission of these frequencies.




Fig. 1. Clockwise rotary spectra for representative moorings and depth levels in the Dardanelles Strait. The major tidal frequencies are marked by dotted lines.




Fig. 2. As in Figure 1, but for the Bosphorus Strait.

Acknowledgments: This work was supported by the Office of Naval Research under Program Element 0601153N.

References

- 1 - Ünlüata Ü, Oguz T., Latif M.A. and Özsoy E., 1990. On the physical oceanography of the Turkish Straits. In: Pratt, L.J. (Ed.), *The Physical Oceanography of Sea Straits*. NATO ASI Ser. Kluwer Academic Publishing, Dordrecht, pp. 25–60.